arXiv:0904.4326v1 [math.DG] 28 Apr 2009

Generalized Nambu dynamics and vectorial Hamiltonians

V.N.Dumachev[1]
dumv@comch.ru

On the basis of Liouville theorem the generalization of the Nambu mechanics is considered. Is shown, that Poisson manifolds of n-dimensional multi-symplectic phase space have inducting by (n-1) Hamiltonian k-vector fields, each of which requires of (k)-hamiltonians.

Keywords: Liouville theorem, Hamiltonian vector fields.

1. Consider a system of the differential equations as submanifold Σ in a jet bundles $J^n(\pi)$: $E \to M$, and [1]

$$F(t, x_0, x_1, ..., x_n) = 0,$$

where $t \in M \subset R$, $u = x_0 \in U \subset R$, $x_i \in J^i(\pi) \subset R^n$, $E = M \times U$. We select from $\Sigma \subset J^n(\pi)$ the equations with Poisson structure and bracket

$$\{H, G\} = X_H \rfloor dG = \mathcal{L}_{X_H} G, \tag{1}$$

and write Cartan distribution as

$$\theta_i = dx_i - \{H, x_i\} dt. \tag{2}$$

Here $\mathcal{L}_{X_H}$ is the Lie derivative along vector field $X_H \in \Lambda^1$, Λ^n a exterior graded algebra of k-vector fields, $H = H(x)$ - while unknown function. Contact vector field $(X_H \rfloor \theta = 0)$ has the form

$$X_H = \frac{\partial}{\partial t} + \{H, x_i\} \frac{\partial}{\partial x_i}. \tag{3}$$

2. Consider classical symplectic mechanics on $J^1(\pi)$. Vector field X_H^1 on symplectic manifold (M, ω) is called Hamiltonian, if the 1-form $\Theta = X_H^1 \rfloor \Omega$ is closed $d\Theta = 0$ and exact (for contractible manifolds) [2]. It allows to find a Hamiltonian H. In this case symplectic form is

$$\Omega = dx_0 \wedge dx_1$$

and

$$\Theta = X_H^1 \rfloor \Omega = dH,$$

where

$$H = \frac{1}{2}\left(x_0^2 + x_1^2\right).$$

Poisson structure (1):

$$\{F, G\} = \frac{\partial F}{\partial x_0} \cdot \frac{\partial G}{\partial x_1} - \frac{\partial F}{\partial x_1} \cdot \frac{\partial G}{\partial x_0}.$$

a inducted by Hamiltonian vector field:

$$X_H^1 = \frac{\partial H}{\partial x_1} \cdot \frac{\partial}{\partial x_0} - \frac{\partial H}{\partial x_0} \cdot \frac{\partial}{\partial x_1}.$$

---

[1] Voronezh Institute of the MVD of the Russia

Cartan distribution defined dynamic Hamiltonian equations in canonical coordinates

$$\frac{\partial x_i}{\partial t} = \{H, x_i\}.$$

Volume of the Cartan differential forms (2)

$$I = \theta_0 \wedge \theta_1 = \Omega - X^1 \rfloor \Omega \wedge dt$$

give as Poincaré integral invariant $i \quad (I = di)$:

$$i = \frac{1}{2}(x_0 dx_1 - x_1 dx_0) - H \wedge dt.$$

According to the Liouville theorem anyone Hamiltonian field conserved of the Volume form, i.e. Lie derivative of the 2-form along vector field X^1_H is zero: $\mathcal{L}_X \Omega = 0$. In other words, generated by a vector fields X^1_H the one-parametrical group symplectic transformations $\{g_t\}$ (phase flow) preserves the canonical two-form Ω, i.e. $g_t \Omega = 0$.

3. Now, we have extended the previous calculations on $J^2(\pi)$. From Liouville theorem we have 3-form

$$\Omega = dx_0 \wedge dx_1 \wedge dx_2$$

volume of the phase space.

Theorem 1. The volume 3-form $\Omega \in \Lambda^3$ are supposed by existence of 2 polyvector Hamiltonian fields: $X^1 \in \Lambda^1$ and $X^2 \in \Lambda^2$.

Proof: On definition

$$\mathcal{L}_X \Omega = X \rfloor d\Omega + d(X \rfloor \Omega) = 0.$$

Since $\Omega \in \Lambda^3$, we see that $d\Omega = 0$, and

$$d(X \rfloor \Omega) = 0.$$

From Poincaré lemma it follows that $X \rfloor \Omega$ is exact, and

$$X \rfloor \Omega = \Theta = dH.$$

1) If $\mathbf{X}^1_H \in \Lambda^1$, then $\Theta \in \Lambda^2$, $H \in \Lambda^1$:

$$X^1_H = \left(\frac{\partial H}{\partial x_1} - \frac{\partial H}{\partial x_2}\right)\frac{\partial}{\partial x_0} + \left(\frac{\partial H}{\partial x_2} - \frac{\partial H}{\partial x_0}\right)\frac{\partial}{\partial x_1} + \left(\frac{\partial H}{\partial x_0} - \frac{\partial H}{\partial x_1}\right)\frac{\partial}{\partial x_2},$$

and

$$\Theta = X^1_H \rfloor \Omega = \sum_{i=0}^{2} dH \wedge dx_i.$$

Poisson bracket $X^1_H \rfloor dG = \{H, G\}$ has the form

$$\{H, G\} = \left(\frac{\partial H}{\partial x_1} - \frac{\partial H}{\partial x_2}\right)\frac{\partial G}{\partial x_0} + \left(\frac{\partial H}{\partial x_2} - \frac{\partial H}{\partial x_0}\right)\frac{\partial G}{\partial x_1} + \left(\frac{\partial H}{\partial x_0} - \frac{\partial H}{\partial x_1}\right)\frac{\partial G}{\partial x_2}$$

2) If $\mathbf{X}_H^2 \in \Lambda^2$, then $\Theta \in \Lambda^1$, $H \in \Lambda^0$,

$$X_H^2 = \frac{1}{2}\left(\frac{\partial H}{\partial x_0} \cdot \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} + \frac{\partial H}{\partial x_1} \cdot \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_0} + \frac{\partial H}{\partial x_2} \cdot \frac{\partial}{\partial x_0} \wedge \frac{\partial}{\partial x_1}\right),$$

and

$$\Theta = X_H^2 \rfloor \Omega = dH.$$

Poisson bracket $X_H^2 \rfloor (dF \wedge dG) = \{H, F, G\}$ has the form

$$\begin{aligned}\{F, G, H\} &= \frac{1}{2}\left[\frac{\partial H}{\partial x_0} \cdot \left(\frac{\partial F}{\partial x_1}\frac{\partial G}{\partial x_2} - \frac{\partial F}{\partial x_2}\frac{\partial G}{\partial x_1}\right)\right. \\ &+ \left.\frac{\partial H}{\partial x_1} \cdot \left(\frac{\partial F}{\partial x_2}\frac{\partial G}{\partial x_0} - \frac{\partial F}{\partial x_0}\frac{\partial G}{\partial x_2}\right) + \frac{\partial H}{\partial x_2} \cdot \left(\frac{\partial F}{\partial x_0}\frac{\partial G}{\partial x_1} - \frac{\partial F}{\partial x_1}\frac{\partial G}{\partial x_0}\right)\right]\end{aligned}$$

4. Let's consider of generalization of the previous calculations on $J^n(\pi)$. Let n-form

$$\Omega = dx_0 \wedge dx_1 \wedge \ldots \wedge dx_{n-1}$$

be Liuville's volume of a phase space.

Theorem 2. Conservation laws for the Liuville's volume $\Omega \in \Lambda^n$ suppose existence of $n-1$ polyvector's Hamilton's fields $X^k \in \Lambda^k \quad (1 \le k \le n-1)$.

Proof: On definition

$$L_X \Omega = X \rfloor d\Omega + d(X \rfloor \Omega) = 0.$$

Since $\Omega \in \Lambda^n$, we see that $d\Omega = 0$, and $d(X \rfloor \Omega) = 0$. From Poincare's lemma it follows that $X \rfloor \Omega$ - is exact, and

$$X \rfloor \Omega = \Theta = dH.$$

If $\mathbf{X}_H^k \in \Lambda^k$, then $\Theta \in \Lambda^{n-k}$, $H \in \Lambda^{n-k-1}$ and $(1 \le k \le n-1)$:

$$X_H^{n-1} = \frac{1}{(n-1)!}\sum_{k=0}^{n-1} \frac{\partial H}{\partial x_k} \cdot \frac{\partial}{\partial x_0} \wedge \frac{\partial}{\partial x_1} \wedge \left[\frac{\partial}{\partial x_k}\right] \wedge \ldots \wedge \frac{\partial}{\partial x_{n-1}},$$

$$X_H^{n-2} = \frac{1}{(n-2)!}\sum_{i<k}^{n-1} \left(\frac{\partial H}{\partial x_i} - \frac{\partial H}{\partial x_k}\right) \cdot \frac{\partial}{\partial x_0} \wedge \left[\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_k}\right] \wedge \ldots \wedge \frac{\partial}{\partial x_{n-1}},$$

$$X_H^{n-3} = \frac{1}{(n-3)!}\sum_{i<k<l}^{n-1} \left(\frac{\partial H}{\partial x_i} - \frac{\partial H}{\partial x_k} + \frac{\partial H}{\partial x_l}\right) \cdot \left[\frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial x_l}\right] \wedge \ldots \wedge \frac{\partial}{\partial x_{n-1}},$$

for which

$$\Theta^1 = X_H^{n-1} \rfloor \Omega = dH,$$

$$\Theta^2 = X_H^{n-2} \rfloor \Omega = \sum_{i}^{n-1} dH \wedge dx_i,$$

$$\Theta^3 = X_H^{n-3} \rfloor \Omega = \sum_{i<j}^{n-1} dH \wedge dx_i \wedge dx_j,$$

$$\Theta^4 = X_H^{n-4} \rfloor \Omega = \sum_{i<j<k}^{n-1} dH \wedge dx_i \wedge dx_j \wedge dx_k,$$

$$\Theta^{n-1} = X_H^{1} \rfloor \Omega = \sum_{i<j<...<k}^{n-1} dH \wedge dx_i \wedge dx_j \wedge ... \wedge dx_k.$$

Proposition. For any n-form Ω and Hamiltonian polyvector field $X_H^k \in \Lambda^k$ existence Poisson Structure $X_H^1 \rfloor dx_i = \{H, x_i\}$, containing k Hamiltonians:

$$X_H^k \rfloor (dF_1 \wedge dF_2 \wedge ... \wedge dF_k) = \{H, F_1, F_2, ..., F_k\}.$$

5. Example. Euler's equations for rigid body in Poisson's form may also be written as

$$\begin{cases} \dot{x} = \quad y - z, \\ \dot{y} = -x + z, \\ \dot{z} = \quad x - y, \end{cases}$$

The vectorial form of this equation we write as

$$\dot{\mathbf{x}} = \mathbf{D}\mathbf{x},$$

where

$$\mathbf{x} = \begin{pmatrix} x \\ y \\ z \end{pmatrix} \quad \text{and} \quad \mathbf{D} = \begin{pmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{pmatrix}.$$

This vector flow is called Hamilton's flow if

$$div\mathbf{D}\mathbf{x} = 0.$$

This implies, that

$$\mathbf{D}\mathbf{x} = rot\mathbf{h}.$$

These expressions are a requirement of a closure of differential form $d\omega = 0:$

$$\omega = (y - x)dy \wedge dz + (-x + z)dz \wedge dx + (x - y)dx \wedge dy = (rot\mathbf{h} \cdot d\mathbf{S}).$$

Using an Homotopy operator we have $\omega = d\nu:$

$$\nu = (\mathbf{h} \cdot d\mathbf{x}) \quad \text{where} \quad \mathbf{h} = \tfrac{1}{3} \begin{pmatrix} y^2 + z^2 - x(y+z) \\ z^2 + x^2 - y(z+x) \\ x^2 + y^2 - z(x+y) \end{pmatrix}, \quad d\mathbf{x} = \begin{pmatrix} dx \\ dy \\ dz \end{pmatrix},$$

or

$$\nu = (y^2 + z^2 - x(y+z))dx + (z^2 + x^2 - y(z+x))dy + (x^2 + y^2 - z(x+y))dz.$$

Hamilton's flow has the form

$$X_h^1 = \left(rot\mathbf{h} \cdot \frac{\partial}{\partial \mathbf{x}} \right) = \left(\frac{\partial h_3}{\partial y} - \frac{\partial h_2}{\partial z} \right) \frac{\partial}{\partial x} + \left(\frac{\partial h_1}{\partial z} - \frac{\partial h_3}{\partial x} \right) \frac{\partial}{\partial y} + \left(\frac{\partial h_2}{\partial x} - \frac{\partial h_1}{\partial y} \right) \frac{\partial}{\partial z},$$

and Poisson bracket $\{H, G\} = X_H \rfloor dG$ gives:

$$\dot{\mathbf{x}} = \{\mathbf{h}, \mathbf{x}\},$$

where $\mathbf{h}$ - vectorial Hamiltonian.

From Lax pair

$$\dot{L} = [ML], \qquad L = \begin{pmatrix} x & z & y \\ z & y & x \\ y & x & z \end{pmatrix}, \qquad M = \frac{1}{2} \begin{pmatrix} 0 & -1 & 1 \\ 1 & 0 & -1 \\ -1 & 1 & 0 \end{pmatrix}$$

for these flow we get two scalar invariants

$$I_1 = trL = x + y + z, \qquad I_2 = \frac{1}{2}\, trL^2 = \frac{3}{2}(x^2 + y^2 + z^2),$$

gives bivectors Hamilton's flow

$$X_I^2 = \frac{1}{2} \left(\frac{\partial I}{\partial x_0} \cdot \frac{\partial}{\partial x_1} \wedge \frac{\partial}{\partial x_2} + \frac{\partial I}{\partial x_1} \cdot \frac{\partial}{\partial x_2} \wedge \frac{\partial}{\partial x_0} + \frac{\partial I}{\partial x_2} \cdot \frac{\partial}{\partial x_0} \wedge \frac{\partial}{\partial x_1} \right).$$

For two Hamiltonians I_1, I_2 it gives $X_{I_1 I_2}^1 = X_{I_1}^2 \rfloor dI_2$:

$$X_{I_1 I_2}^1 = \left(\frac{\partial I_1}{\partial y} \frac{\partial I_2}{\partial z} - \frac{\partial I_1}{\partial z} \frac{\partial I_2}{\partial y} \right) \frac{\partial}{\partial x} + \left(\frac{\partial I_1}{\partial z} \frac{\partial I_2}{\partial x} - \frac{\partial I_1}{\partial x} \frac{\partial I_2}{\partial z} \right) \frac{\partial}{\partial y} + \left(\frac{\partial I_1}{\partial x} \frac{\partial I_2}{\partial y} - \frac{\partial I_1}{\partial y} \frac{\partial I_2}{\partial x} \right) \frac{\partial}{\partial z},$$

and Poisson bracket

$$X_H^2 \rfloor (dF \wedge dG) = \{H, F, G\}$$

with dynamics equations

$$\dot{\mathbf{x}} = \{\mathbf{I}_1, \mathbf{I}_2, \mathbf{x}\}.$$

References

1. Griffiths P., Exterior Differential Systems and the Calculus of Variations, Birkäuser, Boston, 1983.

2. Nambu Y. Generalized Hamiltonian dynamics // Phys.Rev.D, 1973, V.7, N.8, P.5405-5412.